



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: A01N 37/18, A61K 49/00, C12N 5/00, C12N 5/02	A1	(11) International Publication Number: WO 00/19823 (43) International Publication Date: 13 April 2000 (13.04.2000)
--	-----------	--

(21) International Application Number: PCT/US99/23393

(22) International Filing Date: 07 October 1999 (07.10.1999)

(30) Priority Data:

60/103,385 07 October 1998 (07.10.1998) US

60/116,409 19 January 1999 (19.01.1999) US

(60) Parent Application or Grant

THE BOARD OF TRUSTEES OF THE UNIVERSITY OF
ARKANSAS [/]; (). MANOLAGAS, Stavros, C. [/];

(). JILKA, Robert, L. [/]; (). WEINSTEIN, Robert, S. [/];

(). BELLIDO, Teresita [/]; (). ADLER, Benjamin, A. ; ().

Published

(54) Title: METHODS OF SCREENING FOR APOPTOSIS-CONTROLLING AGENTS FOR BONE ANABOLIC THERAPIES AND
USES THEREOF(54) Titre: PROCEDES DE DETECTION D'AGENTS REGULANT L'APOPTOSE DESTINES A LA THERAPIE ANABOLISANTE
OSSEUSE ET UTILISATIONS DE TELS AGENTS

(57) Abstract

The present invention demonstrates that human parathyroid hormone 1-34 [hPTH(1-34)] exerts anti-apoptotic effects on osteoblasts when administered in an intermittent fashion to mice in vivo. The present invention further demonstrates that bovine PTH(1-34) [bPTH(1-34)] prevents glucocorticoid-induced apoptosis of osteoblastic and osteocytic cells in vitro. Therefore, the present invention demonstrates that the previously established anabolic effects of PTH on the skeleton are mediated by its ability to postpone osteoblast apoptosis, as opposed to a stimulatory effect on osteoblastogenesis. The present invention provides methods of screening agents for anti-apoptotic effects on osteoblasts, wherein such agents stimulate and/or restore bone in osteopenic individuals, or prevent bone loss caused by agents such as glucocorticoids.

(57) Abrégé

La présente invention vise à démontrer que l'hormone parathyroïdienne humaine 1-34 [hPTH(1-34)] exerce des effets anti-apoptotiques sur les ostéoblastes lorsqu'on l'administre in vivo de manière intermittente à des souris. La présente invention vise également à démontrer que la PTH(1-34) [bPTH(1-34)] empêche l'apoptose in vivo, induite par les glucocorticoïdes, des cellules ostéoblastiques et ostéocytiques. Cette invention tend par conséquent à démontrer que les effets anabolisants précédemment établis de la PTH sur le squelette résultent de sa capacité à retarder l'apoptose des ostéoblastes, qui s'oppose à un effet stimulant sur l'ostéoblastogenèse. La présente invention se rapporte à des procédés de détection d'agents possédant des pouvoirs anti-apoptotiques sur les ostéoblastes, lesdits agents stimulant et/ou reconstituant l'os chez des sujets ostéopéniques ou empêchant les pertes osseuses au moyen d'agents tels que les glucocorticoïdes.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 37/18, A61K 49/00, C12N 5/00, 5/02	A1	(11) International Publication Number: WO 00/19823 (43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/23393 (22) International Filing Date: 7 October 1999 (07.10.99) (30) Priority Data: 60/103,385 7 October 1998 (07.10.98) US 60/116,409 19 January 1999 (19.01.99) US (71) Applicant: THE BOARD OF TRUSTEES OF THE UNIVERSITY OF ARKANSAS [US/US]; 2404 North University Avenue, Little Rock, AR 72207-3608 (US). (72) Inventors: MANOLAGAS, Stavros, C.; 35 River Ridge Circle, Little Rock, AR 72227 (US). JILKA, Robert, L.; 14202 Clarborne, Little Rock, AR 72211 (US). WEINSTEIN, Robert, S.; 11 Chalmette, Little Rock, AR 72211 (US). BELLIDO, Teresita; 9 Westglen Cove, Little Rock, AR 72211 (US). (74) Agent: ADLER, Benjamin, A.; McGregor & Adler, 8011 Candle Lane, Houston, TX 77071 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHODS OF SCREENING FOR APOPTOSIS-CONTROLLING AGENTS FOR BONE ANABOLIC THERAPIES AND USES THEREOF		
(57) Abstract <p>The present invention demonstrates that human parathyroid hormone 1-34 [hPTH(1-34)] exerts anti-apoptotic effects on osteoblasts when administered in an intermittent fashion to mice <i>in vivo</i>. The present invention further demonstrates that bovine PTH(1-34) [bPTH(1-34)] prevents glucocorticoid-induced apoptosis of osteoblastic and osteocytic cells <i>in vitro</i>. Therefore, the present invention demonstrates that the previously established anabolic effects of PTH on the skeleton are mediated by its ability to postpone osteoblast apoptosis, as opposed to a stimulatory effect on osteoblastogenesis. The present invention provides methods of screening agents for anti-apoptotic effects on osteoblasts, wherein such agents stimulate and/or restore bone in osteopenic individuals, or prevent bone loss caused by agents such as glucocorticoids.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5

10

15

20

25

30

35

40

45

50

55

**METHODS OF SCREENING FOR APOPTOSIS-CONTROLLING AGENTS
FOR BONE ANABOLIC THERAPIES AND USES THEREOF**

BACKGROUND OF THE INVENTION

Cross-Reference to Related Applications

This application claims the benefit of priority of U.S. provisional application 60/116,409, January 19, 1999 and 60/103,385 filed October 7, 1998, now abandoned.

Federal Funding Legend

This invention was produced in part using funds obtained through grant PO1-AG13918 from the National Institutes of Health. Consequently, the federal government has certain rights in this invention.

Field of the Invention

The present invention relates generally to bone physiology. More specifically, the present invention relates to inhibiting apoptosis of osteoblasts and osteocytes.

Description of the Related Art

Remodeling of the human adult skeleton is carried out by teams of juxtaposed osteoclasts and osteoblasts. Osteoclasts and osteoblasts are specialized cell types that originate from hematopoietic and mesenchymal progenitors of the bone marrow, respectively. During bone remodeling, old bone is resorbed by osteoclasts and replaced with new bone by osteoblasts. After they have completed bone matrix synthesis, osteoblasts become osteocytes or lining cells, or they undergo apoptosis.

The osteoblasts and osteoclasts that carry out bone remodeling comprise the basic multi-cellular unit (BMU). Because the lifetime of the basic multi-cellular unit is longer than the lifetime of the individual osteoclasts and osteoblasts, new cells must be continuously supplied from progenitors in the bone marrow for progression to occur. Continuous and orderly supply of these cells, as well as the appropriate rate of apoptosis, is essential for bone homeostasis, as increased or decreased production of osteoclasts or osteoblasts leads to osteoporosis, Paget's, metastatic and renal bone disease. Little is known, however, about the factors that regulate osteogenesis in postnatal life and how osteoblastogenesis and osteoclastogenesis are coordinated to ensure a balance between formation and resorption during remodeling.

During the last few years, it has been established that the process of bone remodeling is regulated locally by growth factors and cytokines produced in the bone micro-environment. In addition, systemic hormones modulate the production and/or action of locally produced cytokines and growth factors, thereby influencing the rate of bone remodeling. Bone morphogenetic proteins (BMPs) are unique among growth factors that influence osteoblast differentiation because

5 they can initiate this process from uncommitted progenitors *in vitro* as well as *in vivo*.

10 Osteoblast commitment is mediated by the type I bone morphogenetic proteins receptor and involves the phosphorylation of
5 specific transactivators (smad 1, 5 and possibly 9), which then oligomerize with smad 4 and translocate into the nucleus. These events induce an osteoblast specific transcription factor (OSF-2/cbfa-
15 1/PEBP2aA/AML3), which in turn activates osteoblast-specific genes (6,7). Bone morphogenetic protein-2 and bone morphogenetic protein-4
10 are expressed during murine embryonal skeletogenesis (day 10-12) and act on cells isolated from murine limb buds to promote their differentiation into osteoblasts. In addition, bone morphogenetic protein-2 and bone morphogenetic protein-4 are involved in fracture
25 healing, as evidenced by their expression in primitive mesenchymal cells and chondrocytes at the site of callus formation, as well as the ability of bone morphogenetic proteins to accelerate the fracture healing
15 process when supplied exogenously.
30

Bone morphogenetic proteins play an essential role in the differentiation of cells that provide support for osteoclast development.
35 20 Osteoclast development requires support from stromal/osteoblastic cells. Moreover, *in vivo*, osteoclastogenesis and osteoblastogenesis proceed simultaneously in most circumstances. This dependency is mediated by a membrane bound cytokine-like molecule
40 (osteoprotegerin ligand/RANK ligand) present in mesenchymal cells which binds to a specific receptor on osteoclast progenitor cells. Such
25 binding is essential, and together with M-CSF, sufficient, for osteoclastogenesis.
45

5 The adverse effects of hypercortisolism on bone have been
recognized for over 60 years, but the precise cellular and molecular
basis of these changes has remained elusive. Today, the iatrogenic form
10 of the disease has become far more common than Cushing's syndrome
5 and glucocorticoid-induced osteoporosis is now third in frequency
following post-menopausal and senile osteoporosis.

15 Bone loss due to glucocorticoid excess is diffuse, affecting
both cortical and cancellous bone, but has a predilection for the axial
skeleton. Spontaneous fractures of the vertebrae or ribs are, therefore,
10 often presenting manifestations of the disorder. A cardinal feature of
20 glucocorticoid-induced osteoporosis is decreased bone formation. In
addition, patients receiving long-term glucocorticoid therapy sometimes
develop collapse of the femoral head (osteonecrosis), but the mechanism
25 underlying this is uncertain. Decreased bone formation, and *in situ*
15 death of isolated segments of the proximal femur suggest that
glucocorticoid excess may alter the birth and death of bone cells.
30 Defective osteoblastogenesis has been reported to be linked to reduced
bone formation and age-related osteopenia in the SAMP6 mouse.
Besides the relationship between aberrant osteoblast production and
20 osteoporosis, it has been recently shown that a significant proportion of
35 osteoblasts undergo apoptosis, which raises the possibility that the
premature or more frequent occurrence of osteoblast apoptosis could
contribute to incomplete repair of resorption cavities and loss of bone.

40 Once osteoblasts have completed their bone-forming
25 function, they either die by apoptosis, become entrapped in bone matrix
and become osteocytes, or remain on the surface as lining cells.
45 Previous studies have demonstrated that the number of osteoblasts is a
critical determinant of bone formation, and that the osteopenic effects

5 of glucocorticoids are due, at least in part, to acceleration of osteoblast apoptosis and stimulation of osteocyte apoptosis.

10 While several agents are capable of decreasing bone resorption and halting further bone loss in osteopenic states, the ideal
5 drug would be an anabolic agent that increases bone mass by rebuilding bone. It is well established that daily injections of low doses of
15 parathyroid hormone (PTH), an agent better known for its role in calcium homeostasis, increases bone mass in animals and humans (5) as
does the PTH-related protein (PTHrP), the only other known ligand of
10 the PTH receptor (6). The mechanism of these anabolic effects,
20 however, has not been established.

The prior art is deficient in methods of inhibiting apoptosis
25 of osteoblasts and osteocytes. The present invention fulfills this long-standing need and desire in the art.

15

35 SUMMARY OF THE INVENTION

20

40 Intermittent PTH administration increases bone mass, but the mechanism of this effect has remained heretofore unknown. Daily PTH injections in mice either with normal bone mass or osteopenia due
45 to defective osteoblastogenesis increased bone formation without
25 affecting the generation of new osteoblasts. Instead, PTH did increase the life span of mature osteoblasts by preventing their apoptosis, an

5 effect reproduced *in vitro*. Increasing the work performed by a cell
population to augment tissue mass by suppressing apoptosis represents
a novel biologic paradigm for regenerating tissues; and could provide a
10 pharmacotherapeutic strategy for rebuilding bone in patients with
5 established osteopenia.

Evidence is presented herein that human parathyroid
hormone 1-34 [hPTH(1-34)] exerts anti-apoptotic effects on osteoblasts
15 when administered in an intermittent fashion to mice *in vivo*. The
present invention also provides evidence that bovine PTH(1-34)
10 [bPTH(1-34)] prevents glucocorticoid-induced apoptosis of osteoblastic
20 and osteocytic cells *in vitro*.

One object of the present invention is to provide methods
for screening compounds that prevent osteoblast apoptosis, thereby
25 stimulating bone formation and/or restoring bone in osteopenic
15 individuals, or preventing bone loss caused by agents such as
glucocorticoids.

30 In an embodiment of the present invention, there is
provided a method of reducing the number of osteoblasts undergoing
apoptosis in an individual in need of such treatment, comprising the
35 20 step of: administering a therapeutic dose of human parathyroid
hormone [hPTH(1-34)] to said individual, wherein administration of
human parathyroid hormone [hPTH(1-34)] results in a reduction in the
40 number of osteoblasts undergoing apoptosis, thereby reducing bone loss
and/or stimulating bone formation in said individual.

25 In another embodiment of the present invention, there is
provided a method of screening compounds that stimulate bone
45 formation, comprising the steps of: (a) contacting osteoblast cells with a
test compound; (b) determining the number of said cells undergoing
50

5 apoptosis; and (c) comparing the number of apoptotic cells with
osteoblast cells that have not been contacted with said compound,
wherein fewer apoptotic cells following contact with said compound
10 than in the absence of said contact indicates that said compound inhibits
5 apoptosis resulting in stimulation of bone formation.

In yet another embodiment of the present invention, there
15 is provided a method of screening for compounds that decrease bone
loss, comprising the steps of: (a) treating osteoblast cells with a
glucocorticoid; (b) contacting said osteoblast cells with a test compound;
10 (c) determining the number of said osteoblast cells undergoing
20 apoptosis; and (d) comparing the number of apoptotic cells with
osteoblast cells that have been treated with said glucocorticoid but were
not contacted with said test compound, wherein fewer apoptotic cells
25 following contact with said test compound than in the absence of said
15 contact with said test compound indicates that said compound inhibits
apoptosis of osteoblast cells thereby reducing bone loss.

30 Other and further aspects, features, and advantages of the
present invention will be apparent from the following description of the
presently preferred embodiments of the invention. These embodiments
35 20 are given for the purpose of disclosure.

40 BRIEF DESCRIPTION OF THE DRAWINGS

45 25 So that the matter in which the above-recited features,
advantages and objects of the invention, as well as others which will
become clear, are attained and can be understood in detail, more
50

5 particular descriptions of the invention briefly summarized above may
be had by reference to certain embodiments thereof which are
10 illustrated in the appended drawings. These drawings form a part of
the specification. It is to be noted, however, that the appended
5 drawings illustrate preferred embodiments of the invention and
therefore are not to be considered limiting in their scope.

15 **Figure 1** shows glucocorticoid-induced apoptosis of
osteoblastic cells inhibited by the specific caspase-3 inhibitor, DEVD.

20 **Figure 2** shows that parathyroid hormone blocks
glucocorticoid-induced, but not TNF α -induced, apoptosis of osteoblastic
cells.

25 **Figure 3** shows that parathyroid hormone blocks
glucocorticoid-induced, but not TNF α -induced, apoptosis of MLO-Y4
osteocytes.

30 **Figure 4** shows that PTH fails to stimulate
osteoblastogenesis.

Figure 5 shows the BMD changes in PTH-treated mice.

35 **Figure 6** shows that PTH stimulates osteoblast and
osteocyte number as well as bone formation rate.

40 **Figure 7** shows that bPTH(1-34) blocks glucocorticoid-
induced apoptosis and bPTH(3-34) prevents the anti-apoptotic effect of
1-34 PTH.

45 **Figure 8** shows that PTH blocks glucocorticoid-induced
apoptosis of osteoblastic cells.

5 **Figure 9** shows that bPTH(1-34) blocks glucocorticoid-induced apoptosis of MLO-Y4 osteocytes and bPTH(3-34) prevents the anti-apoptotic effect of 1-34 PTH.

10 **Figure 10** shows that PTH and the cAMP analog, DBA, block
5 glucocorticoid-induced apoptosis of MLO-Y4 osteocytes.

15 **Figure 11** shows the effect of PTH on BMD. **Figure 11A**, Each point represents the mean (\pm s.d.) change in hindlimb BMD from base line. * $P < 0.05$ vs. vehicle established using a mixed effects longitudinal ANOVA model (Procmixed, SAS, Cary, NC) to allow
20 10 specification of the covariance structure. **Figure 11B**, Mean (\pm s.d.) BMD of hindlimb of SAMR1 and SAMP6 mice prior to ("initial") and after ("final") 28 days of treatment with hPTH(1-34). * $P < 0.05$ vs. initial
25 by paired t-test; $P < 0.05$ vs. SAMR1 by Student's t-test.

Figure 12 shows distal femoral cancellous bone viewed
15 with polarized light to reveal lamellar architecture. Arrows indicate osteocytes. Original magnification = 200X.

30 **Figure 13** shows the mechanism and signal specificity of the suppressive effect of PTH on apoptosis in cultures of osteoblastic and osteocytic cells. **Figure 13A**, Inhibition of dexamethasone-induced
35 20 apoptosis of calvaria cells and MLO-Y4 cells by PTH. Original magnification 400X. Insets: % of cells undergoing apoptosis determined from evaluation of 200 cells in randomly selected fields. **Figure 13B**,
40 Cells (10^4 per cm^2) were incubated for 1 hour in vehicle (Veh) or 10^{-8} M bPTH(1-34), and then for an additional 6 hours in the absence ("basal")
25 or presence of 5×10^{-5} M etoposide ("etop"), 10^{-7} M dexamethasone ("dex"), or 10^{-9} M TNF. **Figure 13C**, Osteoblastic calvaria cells were
45 cultured for 1 hour in vehicle or the indicated log molar concentrations of bPTH(1-34), bPTH(3-34) or DB-cAMP, and then for an additional 6

5 hours in the absence or presence of 10^{-7} M dexamethasone. Adherent
cells were released by digestion with trypsin-EDTA, combined with
nonadherent cells, and apoptotic cells enumerated by trypan blue
10 staining (7). Bars represent the mean (\pm s.d.) of 3 independent
5 measurements. Cell death induced by etoposide, dexamethasone and
TNF was blocked by DEVD-CHO, a cell permeable inhibitor of caspases
required for the execution phase of apoptosis (21). Data were analyzed
15 by ANOVA. Etoposide, dexamethasone, and TNF caused a significant
($p < 0.05$) increase in apoptosis in cultures containing vehicle. * $p < 0.05$
10 vs. vehicle (A), or vs. dexamethasone alone (B).

DETAILED DESCRIPTION OF THE INVENTION

25 Intermittent PTH administration increases bone mass, but
15 the mechanism of this effect has remained heretofore unknown. Daily
PTH injections in mice either with normal bone mass or osteopenia due
30 to defective osteoblastogenesis increased bone formation without
affecting the generation of new osteoblasts. Instead, PTH did increase
35 the life span of mature osteoblasts by preventing their apoptosis, an
20 effect reproduced *in vitro*. Increasing the work performed by a cell
population to augment tissue mass by suppressing apoptosis represents
40 a novel biologic paradigm for regenerating tissues; and could provide a
pharmacotherapeutic strategy for rebuilding bone in patients with
established osteopenia.

25 Evidence is presented herein that human parathyroid
45 hormone 1-34 [hPTH(1-34)] exerts anti-apoptotic effects on osteoblasts
when administered in an intermittent fashion to mice *in vivo*. Evidence

5 is also presented herein that bovine PTH(1-34) [bPTH(1-34)] prevents
glucocorticoid-induced apoptosis of osteoblastic and osteocytic cells *in*
10 *vitro*. These observations demonstrate that the previously established
5 anabolic effects of parathyroid hormone on the skeleton are mediated
by its ability to postpone osteoblast apoptosis, as opposed to a
stimulatory effect on osteoblastogenesis. Results presented herein also
15 demonstrate that the ability of parathyroid hormone to prevent
glucocorticoid-induced osteoblast and osteocyte apoptosis is due to
direct interference with a private death pathway that occurs prior to
10 activation of the final steps of apoptotic mechanism such as activation of
the protease caspase-3.

The present invention is directed towards methods of
screening agents for the ability to inhibit apoptosis of osteoblasts and
25 osteocytes, thereby identifying agents capable of stimulating and/or
restoring bone formation, or preventing bone loss due to treatment with
15 agents such as glucocorticoids.

The present invention is directed to a method of reducing
the number of osteoblasts undergoing apoptosis in an individual in need
of such treatment, comprising the step of: administering a therapeutic
35 20 dose of human parathyroid hormone [hPTH(1-34)] to said individual,
wherein administration of human parathyroid hormone [hPTH(1-34)]
results in a reduction in the number of osteoblasts undergoing
40 apoptosis, thereby preventing bone loss and/or stimulating bone
formation in said individual. In one aspect, the individual is osteopenic.
25 Preferably, the individual is selected from the group consisting of an
individual currently being treated with one or more glucocorticoid
45 compounds and an individual previously treated with one or more
glucocorticoid compounds. Although any route of administration of

5 human parathyroid hormone [hPTH(1-34)] may be used, systemic, oral,
intravenous, nasal spray and inhalation are preferred. Generally, the
human parathyroid hormone [hPTH(1-34)] is administered in a dose of
10 from about 10 µg/kg of body weight to about 1000 µg/kg of body
5 weight.

The present invention is also directed to a method of
15 screening compounds that stimulate bone formation, comprising the
steps of: (a) contacting osteoblast cells with said compound; (b)
determining the number of said cells undergoing apoptosis; and (c)
10 comparing the number of apoptotic cells with osteoblast cells that have
not been contacted with said compound, wherein fewer apoptotic cells
following contact with said compound than in the absence of said
contact indicates that said compound inhibits apoptosis resulting in
25 stimulation of bone formation. Generally, in this method, the

15 contacting of said osteoblast cells is selected from the group
consisting of *in vitro* osteoblast cells and an *in vivo* murine animal
30 model. Representative *in vivo* murine animal models are the SAMP6
mouse and the SAMR1 mouse. Generally, the stimulation of bone
formation is confirmed by methods known to those having ordinary
35 skill in this art such as measuring BMD, measuring cancellous bone area,
measuring cancellous bone formation rate, measuring the number of
osteoblasts per cancellous bone perimeter and measuring the number of
40 osteocytes per bone area in said murine animal model following said
contact with said compound compared with a murine animal model in
25 the absence of said contact with said compound. The determination of
apoptotic cells may be by microscopy of stained cells, TUNEL, Hoescht
45 33258 dye and video image analysis.

5 The present invention is also directed to a method of
screening for compounds that decrease bone loss, comprising the steps
of: (a) treating osteoblast cells with a glucocorticoid; (b) contacting said
10 osteoblast cells with a test compound; (c) determining the number of
5 said osteoblast cells undergoing apoptosis; and (d) comparing the
number of apoptotic cells with osteoblast cells that have been treated
with said glucocorticoid but were not contacted with said test
15 compound, wherein fewer apoptotic cells following contact with said
test compound than in the absence of said contact with said test
20 compound indicates that said compound inhibits apoptosis of osteoblast
cells thereby reducing bone loss. The contacting of the osteoblast cells
may be *in vitro* osteoblast cells or in an *in vivo* murine animal model.
Representative *in vivo* murine animal models include the SAMP6 mouse
25 and the SAMR1 mouse. The determination of apoptotic cells may be by
15 microscopy of stained cells, TUNEL, Hoescht 33258 dye and video image
analysis.

30 It is specifically contemplated that pharmaceutical
compositions may be prepared using the parathyroid hormone of the
present invention. In such a case, the pharmaceutical composition
35 20 comprises the parathyroid hormone of the present invention and a
pharmaceutically acceptable carrier. A person having ordinary skill in
this art would readily be able to determine, without undue
40 experimentation, the appropriate dosages and routes of administration
of this parathyroid hormone of the present invention. When used *in*
25 *in vivo* for therapy, the parathyroid hormone of the present invention is
administered to the patient or an animal in therapeutically effective
45 amounts, i.e., amounts that increase or stimulate bone formation. It will
normally be administered parenterally, preferably subcutaneously by

nasal spray or inhalation, but other routes of administration will be used as appropriate.

The dose and dosage regimen of the parathyroid hormone will depend upon the nature of the disease, the characteristics of the particular parathyroid hormone, *e.g.*, its therapeutic index, the patient, the patient's history and other factors. The amount of parathyroid hormone administered will typically be in the range of about 10 to about 1000 $\mu\text{g/kg}$ of patient weight. The schedule will be continued to optimize effectiveness while balanced against negative effects of treatment. See Remington's Pharmaceutical Science, 17th Ed. (1990) Mark Publishing Co., Easton, Penn.; and *Goodman and Gilman's: The Pharmacological Basis of Therapeutics* 8th Ed (1990) Pergamon Press; which are incorporated herein by reference. For parenteral administration, parathyroid hormone will most typically be formulated in a unit dosage injectable form (solution, suspension, emulsion) in association with a pharmaceutically acceptable parenteral vehicle. Such vehicles are preferably non-toxic and non-therapeutic. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils and ethyl oleate may also be used. Liposomes may be used as carriers. The vehicle may contain minor amounts of additives such as substances that enhance isotonicity and chemical stability, *e.g.*, buffers and preservatives. Parathyroid hormone will typically be formulated in such vehicles at concentrations of about 10 $\mu\text{g/ml}$ to 1000 $\mu\text{g/ml}$.

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion:

EXAMPLE 1*In vitro* effects of PTH

The number of osteoblasts, a critical determinant of bone formation and bone mass, depends both on the birth rate of these cells, which reflects the frequency of cell division of mesenchymal precursors, and on their life span, which reflects the timing of death by apoptosis. *In vivo* evidence indicates that intermittent administration of parathyroid hormone(1-34) increases bone formation and BMD in mice and that these changes are associated with decreased osteoblast and osteocyte apoptosis, but not with increased production of progenitors in the bone marrow.

To determine the mechanism of such actions, the effects of parathyroid hormone(1-34) on the apoptosis of cultured osteoblastic cells isolated from neonatal murine calvaria and the MLO-Y4 osteocyte cell line (provided by L. Bonewald) were examined. Chromatin condensation, nuclear fragmentation, and DNA degradation--cardinal features of apoptotic cells--were monitored by microscopic examination of cells stained with the DNA dye Hoescht 33258, or stably transfected with green fluorescent protein gene containing a nuclear localization sequence, and by DNA end labeling (TUNEL). Enumeration of apoptotic cells was performed by trypan blue staining, and correlated closely with morphologic changes and TUNEL.

In both osteoblast and osteocyte cultures, 15% of the cells were apoptotic 6 hours after addition of 100 nM dexamethasone, as compared to 4% in cultures without the steroid. This effect was completely prevented by 10 nM parathyroid hormone(1-34) or 1 mM dibutyryl-cAMP added 1 hour prior to addition of dexamethasone. The parathyroid hormone effect did not involve cytokines with anti-

5 apoptotic properties, as neutralizing antibodies against IL-6, IL-11 or
LIF did not interfere with this phenomenon. In contrast to
dexamethasone-induced apoptosis, parathyroid hormone had no
10 influence on TNF-induced apoptosis. These findings are consistent with
5 *in vivo* evidence demonstrating that the anabolic effects of parathyroid
hormone are due to its anti-apoptotic effects on osteoblasts and
osteocytes; and that parathyroid hormone interferes directly with a
15 private apoptosis pathway at a site(s) upstream of the induction of the
degradation phase of apoptosis which is executed by caspase-3.

10

20 EXAMPLE 2

Calvaria and MLO-Y4 cells

Osteoblastic calvaria cells (9) were cultured in α MEM (Gibco-
25 BRL, Grand Island, NY) supplemented with 10% FBS (Sigma Chemical Co.,
15 St. Louis, MO). Murine osteocyte-like MLO-Y4 cells stably transfected
with EGFP were cultured on collagen coated plates in MEM
30 supplemented with 5% FBS and 5% bovine calf serum. Cultures were
maintained for 6 hours in the presence of 10^{-7} M dexamethasone
without or with preincubation for 1 hour with 10^{-8} M bPTH(1-34) and
20 fixed in neutral buffered formalin. The pyknotic fragmented nuclei
35 (arrows) typical of apoptotic cells were visualized with Hoescht 33258
fluorescent dye (Polysciences, Inc., Bayshore, NY), used at a
concentration of 1 μ g/ml in 0.5 M NaCl, 10 mM Tris-HCl, 1 mM EDTA, pH
40 7.4) in osteoblastic calvaria cells, and by EGFP fluorescence in MLO-Y4
25 osteocytes.

Osteoblastic cells were isolated from calvaria of 3- to 6-day-
45 old C57/B1 mice by sequential collagenase digestion. Cells were cultured
for 5-8 days in α MEM supplemented with 10% FBS and frozen in liquid

5 N₂ until use. MLO-Y4 cells (provided by Dr. L. Bonewald, University of
Texas Health Science Center at San Antonio, San Antonio, TX) were
transduced with the pLXSN retroviral vector containing a construct
10 encoding enhanced green fluorescent protein (Clontech, Palo Alto, CA)
5 with the SV40 large T antigen nuclear localization sequence [D. Kalderon
et al., Cell 39, 499 (1984)] attached to the carboxyterminus. Stably
transduced cells were selected for neomycin resistance using G418
15 (Sigma, St. Louis, MO).

10 EXAMPLE 3

20 Mice

4-5 month old male or female SAMR1 and SAMP6 were
given daily injections of vehicle (0.9% saline, 0.01 mM β -
25 mercaptoethanol, 0.1 mM acetic acid) or 400 ng/g body weight of
15 hPTH(1-34) (Bachem, Torrance, CA) dissolved in vehicle (n=6-7 per
group). Mice were fed a standard rodent diet (Agway RMH 3000,
Arlington Heights, IL) *ad libitum*. The BMD of the spine and
hindquarters was determined one day prior to initiation of the
experiment (baseline scan) and at weekly intervals thereafter using
30 dual-energy X-ray absorptiometry (QDR 2000, Hologic, Inc.) as
20 described previously (3). The evaluation of each scan was based on the
exact positioning and region of interest placement of the baseline scan
using the "Compare" technique (4).
40

25 EXAMPLE 4

Experimental Methods

45 To determine the effect of PTH on osteoblast formation,
function and fate, mice from the experiment shown in Figure 11 were
50

5 killed on day 28. The animals had been pretreated with tetracycline (5
µg/g, s.c.) on day 19 and 26. Osteoblast progenitors were measured
10 using marrow cells from one femur. Cells from each animal were
cultured separately at 2.5×10^6 per 10 cm² well and maintained for 28
5 days in phenol red-free (MEM containing 15% preselected FBS (HyClone,
Logan, UT) and 1 mM ascorbate-2-phosphate (18). Von Kossa's method
15 was used to visualize and enumerate colonies containing mineralized
bone matrix. Because each colony is derived from a single osteoblast
progenitor, the colony forming unit osteoblast (CFU-OB), the number of
20 CFU-OB colonies reflects the number of osteoblast progenitors present in
the original bone marrow isolate. The remaining femur and lumbar
vertebrae were fixed in 4°C Millonig's phosphate-buffered 10%
formalin, pH 7.4 and embedded undecalcified in methyl methacrylate.

25 Measurements of the femoral length and the midshaft
15 diaphyseal cortical width were made with a digital caliper at a
resolution of 0.01 mm (Mitutoyo Model #500-196, Ace Tools, Ft. Smith,
30 AR). Histomorphometric examination of five micron thick bone sections
were performed using a computer and digitizer tablet (OsteoMetrics Inc.
Version 3.00, Atlanta, GA) interfaced to a Zeiss Axioscope (Carl Zeiss,
35 20 Inc., Thornwood, NY) with a drawing tube attachment (4).
Measurements were confined to the secondary spongiosa of the distal
femur. The terminology and units used are those recommended by the
40 Histomorphometry Nomenclature Committee of the American Society
for Bone and Mineral Research (19). The rate of bone formation
25 ($\mu\text{m}^2/\mu\text{m}/\text{d}$) was calculated from the extent of bone surface labeled with
tetracycline (visualized by fluorescence under UV illumination) and the
45 distance between the labels in areas where two labels are present.

5 Osteoid was recognized by its distinct staining characteristics and osteoblasts were identified as plump cuboidal cells on osteoid surfaces.

10 Apoptotic osteoblasts were detected in sections of nondecified vertebral bone by the TUNEL reaction (TdT-mediated
5 dUTP nick end labeling) using reagents from Oncogene (Cambridge, MA) (4,7). Briefly, sections were incubated in 0.5% pepsin in 0.1 N HCl for 20 minutes at 37°C, rinsed with TBS and then incubated in 30% H₂O₂ in
15 methanol for 5 minutes, and rinsed again. To improve the sensitivity of the reaction, sections were subsequently incubated for 1-2 minutes
10 with 0.15% CuSO₄ in 0.9% NaCl (20). TUNEL-positive hypertrophic chondrocytes were observed at the bottom of the growth plates serving
20 as an internal positive control for each bone section. The prevalence of osteoblast apoptosis was determined by inspecting 1190 and 852
25 osteoblasts in sections from vehicle-treated and 2514 and 1490
15 osteoblasts in PTH-treated SAMR1 and SAMP6 mice, respectively. For the determination of osteocyte apoptosis, 1579 and 1714 osteocytes
30 were evaluated in vehicle-treated and 2930 and 2259 osteocytes in PTH-treated SAMR1 and SAMP6 mice, respectively. Wall width
35 represents the amount of bone synthesized by a team of osteoblasts and
20 was measured as the distance from the bone surface to a cement line in the underlying bone demarcating the site at which bone formation
40 began. Osteocytes were identified inside lacunae in mineralized bone. Osteoclasts were recognized by staining with tartrate resistant acid
phosphatase.

25 The decrease in the number of apoptotic osteoblasts in PTH-treated mice was confirmed in a second set of bone sections stained
45 with the standard TUNEL technique (4,7), as opposed to the modified TUNEL method used for the data presented in the text and Table 1, in

5 which CuSO_4 is added subsequent to peroxidase staining to intensify the
chromogen so as to allow detection of DNA degradation as early as
possible (20). Using the standard TUNEL technique, the prevalence of
10 apoptotic osteoblasts in PTH-treated mice was 0.017% (1 out of 5,900
5 osteoblasts inspected in 36 sections taken from 12 mice: 6 SAMR1 and
6 SAMP6) compared to 0.37% in the vehicle-treated controls (12 out of
3,250 osteoblasts inspected in 39 sections taken from 13 mice, 6 SAMR1
15 and 7 SAMP6); $p < 0.001$ vs. PTH-treated animals by logistic regression
(LogXact, Cytel Corp. Cambridge, MA). The increase in the absolute
10 number of TUNEL-labeled cells observed with the modified as compared
20 to the standard technique is consistent with the contention that the
former procedure allows visualization of cells undergoing the DNA
degradation phase at an earlier stage. Prolongation or shortening of the
25 time that apoptosis can be observed in a specimen as a result of using a
15 more or less sensitive detection method must be taken into
consideration when comparing estimates of the prevalence of the
phenomenon in different studies (4,7). This variability of the duration
30 of the apoptosis process in different cell types can account for reports
estimating the TUNEL-labeled phase of apoptosis from as little as 1.5 to
20 as much as 48 hours.

EXAMPLE 5

In vivo effects on mice of bPTH

25 Intermittent administration of parathyroid hormone exerts
an anabolic effect on the skeleton of animals and humans, most likely
45 due to an increase in the number of osteoblasts. This number depends
both on the birth rate (reflecting the frequency of division of

mesenchymal progenitors) and the life span (reflecting the timing of apoptosis). The present invention examines the effects of intermittent parathyroid hormone administration in mice with either normal (SAMR1) or defective (SAMP6) osteoblastogenesis at four months of age, a time at which both strains have achieved peak bone mass.

Mice (6-7 per group) were given daily subcutaneous injections of 400 ng bovine parathyroid hormone (1-34) per gram of body weight or vehicle for a period of 4 weeks. BMD was monitored weekly by DEXA. One femur was used for tetracycline based dynamic histomorphometry and the other for determination of osteoblast progenitors in *ex vivo* bone marrow cultures. Spine and hindquarter BMD increased gradually in parathyroid hormone-treated mice of either strain reaching 4% and 15%, respectively, over the pretreatment values by 4 weeks. Parathyroid hormone also increased cancellous bone area and bone formation rate (2-3 fold), as well as the number of osteoblasts per cancellous bone perimeter and the number of osteocytes per bone area (2-fold) in both strains.

These changes could not be accounted for by hypertrophy of lining cells and their subsequent conversion to osteocytes because the increased osteocyte density in the expanded cancellous bone area was much too great. Unlike the increase in mature cell numbers, the number of CFU-OB formed per 10^6 marrow cells in *ex vivo* bone marrow cultures did not change in either strain (SAMR1: vehicle= 34 ± 7 , PTH= 34 ± 9 ; SAMP6: vehicle= 17 ± 2 , PTH= 21 ± 5). Parathyroid hormone did, however, decrease osteoblast apoptosis.

As detected by DNA end-labeling (TUNEL), 0.33% and 0.40% of osteoblasts were apoptotic in bone of SAMR1 and SAMP6 mice, respectively (N=3250), whereas only 0.02% of the 4200 osteoblasts

5 examined in parathyroid hormone-treated mice were apoptotic ($P < 0.02$
by z-test). Based on an osteoblast lifetime of 300 hours, and a 3 hour
duration of apoptosis, it was calculated that parathyroid hormone
10 postponed apoptosis of the 30-40% of osteoblasts that would normally
5 undergo this process during bone remodeling. These findings indicate
that suppression of osteoblast apoptosis, rather than increased
osteoblastogenesis, is the mechanism by which intermittent
15 administration of parathyroid hormone stimulates bone formation. This
effect is sufficient to account for the increase in the number of
10 osteoblast and osteocytes and, thereby, the anabolic effect of the
20 hormone.

EXAMPLE 6

In vivo effects of hPTH on mice

25 15 Daily subcutaneous injections of hPTH (1-34) over a four
week period progressively increased bone mineral density (BMD) in
adult mice with normal bone mass (SAMR1) or in mice with osteopenia
30 (3) due to impaired osteoblastogenesis and decreased bone formation
(SAMP6) (Figure 11A). This effect was greater in the hindlimbs than in
20 the spine and did not involve bone growth as reflected by the lack of
35 change in the length of the femur or its width at the diaphysis (Table 1).
Remarkably, the increase in BMD was similar in the two strains, even
though the baseline values were different (Figure 11B).

40 25 Consistent with the BMD increase, histomorphometric
analysis of sections of cancellous bone from the distal femurs of mice of
either strain treated with PTH showed increased cancellous bone area
45 that correlated with the increase in hindlimb BMD ($r = 0.53$, $P < 0.001$).
The latter could be explained by the increase in the number of

osteoblasts covering the bone surface and by the increased rate of bone formation (Table 1). As expected from these changes, the amount of osteoid, the matrix produced by osteoblasts, was also increased, as was the amount of bone estimated to be produced by each team of osteoblasts (wall width). The number of osteoclasts on the cancellous bone surface was not changed with PTH treatment, indicating that bone resorption and turnover were not affected by the hormone. The newly formed bone had normal lamellar architecture as opposed to woven bone seen in severe hyperparathyroidism (Figure 12).

As osteoblasts are short-lived cells (approximately 200 hours in mice) (4,7), the increase in the number of osteoblasts seen in the PTH-treated mice could result from either an increase in the formation of new osteoblasts or the prolongation of their life span. Enumeration of osteoblast progenitors in ex vivo bone marrow cell cultures, a reliable index of *de novo* osteoblastogenesis (3), showed no difference between PTH and vehicle-treated animals (Table 1). The lack of effect of PTH on osteoblast progenitors was documented in the normal mouse strain (SAMR1) as well as in the strain with the diminished baseline osteoblastogenesis (SAMP6), clearly demonstrating that an increase in the formation of new osteoblasts could not account for the increased osteoblast numbers seen in both strains.

Strikingly, however, the percentage of osteoblasts undergoing apoptosis, as determined by TUNEL-labeling, was greatly decreased in animals of either strain receiving PTH (Table 1). Specifically, the average number of apoptotic osteoblasts, determined separately for each animal, was $0.4 \pm 0.5\%$ (range 0-1.4%, n=6) for PTH-treated SAMR1 as compared to $1.7 \pm 1.0\%$ (range 0.5-3.2%, n=6) for vehicle-treated SAMR1, $p < 0.05$. Similarly, the average number of

apoptotic osteoblasts in PTH-treated SAMP6 was $0.1 \pm 0.3\%$ (range 0-0.8%, n=6) as compared to $2.2 \pm 1.4\%$ (range 0.9-4.7%, n=7) for vehicle-treated SAMP6, $p < 0.05$ (8).

Table 1 shows the effect of PTH on osteoblast formation, function and fate. Mice from the experiment shown in Fig. 1 were killed on day 28. The animals had been pretreated with tetracycline (5 $\mu\text{g/g}$, s.c.) on day 19 and 26. Osteoblast progenitors were measured using marrow cells from one femur. The average number of nucleated cells obtained from the femur of PTH treated animals ($19.8 \pm 3.2 \times 10^6$ from SAMR1; $24.3 \pm 3.0 \times 10^6$ from SAMP6) was indistinguishable from animals receiving vehicle ($21.0 \pm 2.7 \times 10^6$ from SAMR1; $21.5 \pm 3.2 \times 10^6$ from SAMP6). The remaining femur and lumbar vertebrae were fixed and embedded undecalcified in methylmethacrylate (3,4,15). Femurs were used for histomorphometric analysis and vertebrae were used for apoptosis determinations. Because osteoblasts in remodeling bone comprise a team, they were identified as cuboidal cells in a row of at least three, lining the osteoid-covered trabecular perimeter. Osteocytes were identified inside lacunae of mineralized cancellous bone. For detection of apoptotic cells, sections were incubated with CuSO_4 to enhance staining of the peroxidase reaction product during the TUNEL procedure, as described in Methods. Osteoblasts and osteocytes exhibiting both brown staining due to TUNEL and the morphological feature of nuclear condensation were counted as apoptotic. With these precautions, TUNEL has been unequivocally associated with apoptosis of osteoblasts and osteocytes in bone (4,29,50). See Table 2 for a summary of apoptotic and non-apoptotic cell counts in individual animals. The data shown represent the mean (\pm s.d.) of each measurement determined from bones from each animal.

5 ^aP<0.05 vs. vehicle by one-tailed Students t-test. ^bTwo-way ANOVA was
used to detect overall effects of PTH. ^cP<0.001 vs. vehicle by GSK
categorical general linear regression (26).

10 Table 2 shows the effect of PTH on apoptosis of osteoblasts
5 and osteocytes in vertebral cancellous bone. Osteoblasts (OB) and
osteocytes (OCT) were identified in sections of lumbar vertebrae, and
those exhibiting both brown staining due to TUNEL and pyknotic nuclei
15 were counted as apoptotic as described in Table 1. Results are from two
separate TUNEL staining procedures. In the first, TUNEL was performed
10 without CuSO₄ enhancement ("w/o Cu"), and data were pooled from
animals from each group for statistical analysis because of the low
20 number of apoptotic osteoblasts visualized with this method. In the
second, CuSO₄ ("with Cu") was used to enhance TUNEL staining, and
25 counts from each animal are shown. With the exception of the apoptotic
15 osteocyte counts in the vehicle treated SAMP6 group, there were no
significant differences among the animals within each group. Pooled
30 data ("w/o Cu") were analyzed by logistic regression. ^aP<0.0001 vs.
vehicle. Data from individual animals were analyzed by GSK. ^bP<0.001
vs. vehicle.

Table 1.

		SAMR1		SAM
		vehicle	PTH	vehicle
5				
10	CFU-OB (# per 10 ⁶ marrow cells)	34 ± 7	34 ± 9	17 ± 3
15	Osteoblast perimeter (%)	7.1 ± 2.9	11.8 ± 6.8 ^A	5.7 ± 3.5
20	Apoptotic osteoblasts (%)	1.7 ± 1.0	0.4 ± 0.5 ^C	2.2 ± 1.4
25	Bone area (% of tissue area)	9.0 ± 4.2	23.2 ± 11.3 ^A	8.9 ± 2.9
30	Bone formation rate (μm/μm ² /d)	0.087 ± 0.039	0.361 ± 0.413 ^A	0.071 ± 0.010
35	Mineralizing perimeter (%)	8.92 ± 4.67	14.96 ± 4.82 ^A	6.56 ± 3.24
40	Mineral apposition rate (μm/d)	1.09 ± 0.43	1.14 ± 0.87	1.07 ± 0.23
45	Trabecular width (μm)	37.2 ± 11.0	56.7 ± 21.2 ^A	39.9 ± 9.1
50	Wall width (μm)	10.2 ± 2.5	13.4 ± 3.2 ^A	5.4 ± 0.9
55	Osteoid perimeter (%)	13.2 ± 7.8	17.4 ± 4.9	9.9 ± 5.5
	Osteocyte areal density (# per bone area)	8.2 ± 3.1	14.8 ± 6.4 ^A	14.3 ± 5.7
	Apoptotic osteocytes (%)	1.7 ± 0.5	0.2 ± 0.2 ^C	2.5 ± 2.0
	Osteoclast perimeter (%)	1.3 ± 0.8	0.7 ± 0.8	2.3 ± 2.5
	Femoral length (mm)	15.82 ± 0.51	16.14 ± 0.48	15.03 ± 0.66
	Diaphyseal cortical width (mm)	0.59 ± 0.38	0.54 ± 0.13	0.56 ± 0.13

Table 2.

Group	Sample, or Mouse ID	Apop totic OB	Total OB Counte d	% Apop totic OB	Apop totic OCT	Total OCT Count ed	% Apopt otic OCT
SAMR1, Veh	Pooled (w/o Cu)	5	1500	0.3	-	-	-
	5-6 (with Cu)	5	271	1.8	7	368	1.9
	5-15	9	280	3.2	9	350	2.5
	5-17	6	258	2.3	4	240	1.6
	5-20	1	199	0.5	2	237	0.8
	5-14	1	113	0.9	4	244	1.6
	5-1	1	69	1.4	2	98	1.8
			Mean \pm s.d:	1.7 \pm 1.0			1.7 \pm 0.5
SAMR1, PTH	Pooled (w/o Cu)	0	3700	0 ^a	-	-	-
	5-7 (with Cu)	0	206	0	0	403	0
	5-8	1	328	0.3	1	477	0.3
	5-18	1	372	0.3	2	489	0.3
	5-2	3	206	1.4	2	618	1.4
	5-9	2	397	0.5	1	669	0.5
	5-11	0	151	0	0	262	0
			Mean \pm s.d:	0.4 \pm 0.5 ^a			0.2 \pm 0.2 ^a

5

10	SAMP6, Veh	Pooled (w/o Cu)	7	1750	0.4	-	-	-
		7-19 (with Cu)	6	180	3.4	9	204	4.3
		7-11	3	64	4.7	4	302	1.3
15		7-6	1	114	0.9	10	169	5.7
		7-12	3	185	1.6	1	353	0.3
		7-17	2	109	1.8	9	342	2.6
		7-21	1	88	1.1	1	208	0.5
20		7-24	2	113	1.8	11	384	2.8
				Mean \pm s.d:	2.2 \pm 1.4			2.5 \pm 2.0
25	SAMP6, PTH	Pooled (w/o Cu)	1	2200	0.05 ^a	-	-	-
		7-1 (with Cu)	0	289	0	3	500	0.6
		7-4	0	387	0	1	487	0.2
		7-7	3	400	0.8	2	350	0.7
30		7-9	0	160	0	2	321	0.6
		7-16	0	120	0	1	293	0.3
		7-23	0	134	0	0	303	0
				Mean \pm s.d:	0.1 \pm 0.3 ^b			0.4 \pm 0.3 ^b

35

40

45

50

55

EXAMPLE 7Effect of PTH on osteocytes

PTH-treated mice also exhibited increased osteocyte density - i.e. number per cancellous bone area (Table 1 and Figure 12). Osteocytes are former osteoblasts that have completed their bone-forming function and are encased within lacunae of the mineralized bone matrix, one of the three possible fates of matrix synthesizing cells, the other two being apoptosis and conversion to lining cells. Hence, an increase in osteocyte density is consistent with, and can only be accounted for by, a suppression of osteoblast apoptosis. Besides the effect on osteoblast apoptosis, intermittent PTH administration also inhibited osteocyte apoptosis (Table 1).

EXAMPLE 8Anti-apoptotic effect of PTH

To determine whether the anti-apoptotic effect of PTH was due to direct action of the hormone on osteoblasts and osteocytes, as opposed to indirect actions mediated by compensatory changes, the effect of PTH on apoptosis was examined using cell cultures. Induction of apoptosis by either etoposide or dexamethasone was attenuated by PTH in primary cultures of osteoblasts isolated from neonatal murine calvaria (9) whereas induction of apoptosis by TNF was not (Figure 13A,B). These findings were reproduced using murine osteoblastic MC3T3-E1 cells as well as human osteoblastic MG-63 cells (10). The anti-apoptotic effect of PTH could be blocked by the PTH/PTHrP receptor antagonist bPTH(3-34) and was mimicked by dibutyryl-cAMP, indicating that it was mediated through the PTH/PTHrP receptor and

subsequent activation of adenylate cyclase (Figure 13C). PTH also prevented etoposide- and dexamethasone-induced apoptosis, but not TNF-induced apoptosis in an established murine osteocyte-like cell line, MLO-Y4 (11), stably transfected with the enhanced green fluorescent protein (EGFP) gene containing a nuclear localization sequence, (Figure 13A,B).

Summary

The data presented herein demonstrates that intermittent administration of PTH stimulates bone formation, not by increasing the proliferation of osteoblast precursors but by preventing osteoblast apoptosis - the fate of the majority of these cells under normal conditions (4, 7) - thereby prolonging the time spent in performing their matrix synthesizing function. The anti-apoptotic effect of PTH is exerted directly on osteoblasts, requires binding of the hormone to the PTH/PTHrP receptor, is mediated by cAMP-generated signals that interfere with some but not all death pathways, and occurs upstream of the common executing phase of apoptosis. Consistent with the results of the present studies in mice, the ability of intermittent administration of PTH to increase osteoblast numbers was not accompanied by an increase in the replication of osteoblast progenitors in the rat (12). Moreover, the demonstration of the ability of PTH to inhibit osteoblast apoptosis is in full agreement with the anti-apoptotic effect of PTHrP on chondrocytes during endochondral bone development (13).

Results of a clinical study have shown that daily subcutaneous injections of PTH is an effective treatment for glucocorticoid-induced osteoporosis (15). The decreased bone formation rate and wall thickness of trabeculae, indicators of diminished work by

5 osteoblasts and the *in situ* death of portions of bone that characterize
glucocorticoid-induced osteoporosis can be accounted for by a
suppressive effect of glucocorticoids on osteoblastogenesis and
10 promotion of apoptosis of osteoblasts and osteocytes (4). The
5 elucidation of the anti-apoptotic effects of PTH *in vivo* along with the
evidence that PTH antagonizes the pro-apoptotic effects of
glucocorticoids *in vitro* are in full agreement with the published clinical
15 observations. Furthermore, they provide both a mechanistic
explanation for the efficacy of PTH in glucocorticoid-induced
10 osteoporosis as well as compelling evidence that its anti-apoptotic
properties make PTH a rational pharmacotherapeutic choice for this
20 condition.

Apart from decreased cell death, another potential source of
25 new osteoblasts is the lining cells that cover quiescent bone surfaces
15 (16). These cells were once matrix synthesizing osteoblasts, and have
escaped apoptosis or encasement within bone as osteocytes to remain
on the bone surface. It has been suggested that PTH can stimulate
30 lining cells to undergo hypertrophy and to resume matrix synthesis
(12,17). Such hypertrophy was not observed, but a contribution from
20 this mechanism cannot be excluded from the results of the present
report. Nevertheless, the magnitude of the anti-apoptotic effect of PTH
35 makes it unnecessary to invoke another explanation for increased bone
formation.

40 In conclusion, the data presented herein demonstrate that
25 prevention of osteoblast apoptosis is the principal mechanism for the
anabolic effects of PTH on bone. Increasing the work-output of a cell
population by suppressing apoptosis represents a novel biologic
45 paradigm for regenerating tissues in general and a rational

5 pharmacotherapeutic strategy for rebuilding bone in particular. PTH,
and possibly PTH mimetics and non-peptide inhibitors of private
apoptotic pathways in osteoblasts, should provide much needed
10 therapies for osteopenias, in particular those in which osteoblast
5 progenitors are low, such as age-related and glucocorticoid induced
osteoporosis.

The following references were cited herein:

- 15 1. S.C. Manolagas & R.L. Jilka, N.Engl.J.Med. 332, 305 (1995).
2. R.L. Jilka et al., Science 257, 88 (1992).
- 10 3. R.L. Jilka et al., J.Clin.Invest. 97, 1732 (1996).
- 20 4. R.S. Weinstein et al., ibid., 102, 274 (1998).
5. D.W. Dempster et al., Endocr.Rev. 14, 690 (1993); J. Reeve et al.,
Lancet 1(7968), 1035 (1976); J. Reeve et al., Br.Med.J. 280, 1340
25 (1980); C.S. Tam et al., Endocrinology 110, 506 (1982); J.S. Finkelstein et
15 al., JAMA 280, 1067 (1998).
6. A.F. Stewart, Bone 19, 303 (1996); B.H. Vickery et al., J. Bone
30 Miner. Res. 11, 1943 (1996).
7. R.L. Jilka et al., J.Bone Miner.Res. 13, 793 (1998).
8. W. Bursch et al., Carcinogenesis 11, 847 (1990); M. Pompeiano et
20 al., Cell Death Differ. 5, 702 (1998).
- 35 9. R.L. Jilka, Bone. 7, 29 (1986).
10. H. Sudo et al., J.Cell Biol. 96, 191 (1983); R.T. Franceschi et al.,
J.Biol.Chem. 263, 18938 (1988).
- 40 11. Y. Kato et al., J.Bone Miner.Res. 12, 2014 (1997).
- 25 12. H. Dobnig & R.T. Turner, Endocrinology 136, 3632 (1995).
13. M. Amling et al, J.Cell Biol. 136, 205 (1997); K. Lee et al.,
45 Endocrinology 137, 5109 (1996).
14. E.M. Aarden et al., J. Cellular Biochem. 55, 287 (1994).

- 5 15. N.E. Lane et al., J.Clin.Invest. 102, 1633 (1998).
16. S.C. Miller & W.S.S. Jee, Bone. Volume 4: Bone Metabolism and Mineralization, B.K. Hall, Ed. (CRC Press, Boca Raton, 1992), p. 1.
- 10 17. D. Leaffer et al., Endocrinology 136, 3624 (1995).
- 5 18. R.L. Jilka et al., J.Clin.Invest. 101, 1942 (1998).
19. A.M. Parfitt et al., J.Bone Miner.Res. 2, 595 (1987).
20. S. Hsu & E. Soban, J. Histochem. Cytochem. 30, 1079 (1982); B.G. Short et al., ibid, 45, 1299 (1997).
- 15 21. S.C. Wright et al., J.Exp.Med. 186, 1107 (1997); N.A. Thornberry & Y. Lazebnik, Science 281, 1312 (1998).
- 10

20 Any patents or publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. Further, these patents and publications are incorporated by reference herein to the same extent as if each

25 individual publication was specifically and individually indicated to be incorporated by reference.

15

30 One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the

35 methods, procedures, treatments, molecules, and specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to

40 those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

25

Claims

5

10

15

20

25

30

35

40

45

50

55

WHAT IS CLAIMED IS:

1. A method of reducing the number of osteoblasts undergoing apoptosis in an individual in need of such treatment, comprising the step of:

administering a therapeutic dose of human parathyroid hormone [hPTH(1-34)] to said individual, wherein administration of human parathyroid hormone [hPTH(1-34)] results in a reduction in the number of osteoblasts undergoing apoptosis, thereby preventing bone loss and/or stimulating bone formation in said individual.

2. The method of claim 1, wherein said individual is osteopenic.

3. The method of claim 1, wherein said individual is selected from the group consisting of an individual currently being treated with one or more glucocorticoid compounds and an individual previously treated with one or more glucocorticoid compounds.

4. The method of claim 1, wherein said administration is selected from the group consisting of systemic, oral, intravenous, nasal spray and inhalation.

5. The method of claim 1, wherein said human parathyroid hormone [hPTH(1-34)] is administered in a dose of from about 10 µg/kg to about 1000 µg/kg.

5

6. A method of screening for a compound that stimulates bone formation, comprising the steps of:

10

(a) contacting osteoblast cells with a test compound;

5

(b) determining the number of said cells undergoing apoptosis; and

15

(c) comparing the number of apoptotic cells with osteoblast cells that have not been contacted with said compound, wherein fewer apoptotic cells following contact with said compound than in the absence of said contact indicates that said compound inhibits apoptosis resulting in stimulation of bone formation.

20

10

25

7. The method of claim 6, wherein said contacting of said osteoblast cells is selected from the group consisting of *in vitro* osteoblast cells and an *in vivo* murine animal model.

15

30

35

8. The method of claim 7, wherein said *in vivo* murine animal model is selected from the group consisting of a SAMP6 mouse, a SAMR1 mouse and other strains of mice.

20

40

9. The method of claim 6, wherein said stimulation of bone formation is confirmed by methods selected from the group consisting of measuring BMD, measuring cancellous bone area, measuring cancellous bone formation rate, measuring the number of osteoblasts per cancellous bone perimeter and measuring the number of osteocytes per bone area in said murine animal model following said

45

25

50

55

5 contact with said compound compared with a murine animal model in
the absence of said contact with said compound.

10 10. The method of claim 6, wherein said determination of
5 apoptotic cells is selected from the group consisting of microscopy of
stained cells, TUNEL, Hoescht 33258 dye and video image analysis.

15 11. A method of screening for a compound that reduces
bone loss, comprising the steps of:

- 20 10 (a) treating osteoblast cells with a glucocorticoid;
(b) contacting said osteoblast cells with a test compound;
25 (c) determining the number of said osteoblast cells
undergoing apoptosis; and
(d) comparing the number of apoptotic cells with
30 15 osteoblast cells that have been treated with said glucocorticoid but were
not contacted with said test compound, wherein fewer apoptotic cells
following contact with said test compound than in the absence of said
35 contact with said test compound indicates that said compound inhibits
apoptosis of osteoblast cells thereby reducing bone loss.

20

40 12. The method of claim 11, wherein said contacting of
said osteoblast cells is selected from the group consisting of *in vitro*
osteoblast cells and an *in vivo* murine animal model.

45

50

55

5

13. The method of claim 12, wherein said *in vivo* murine animal model is selected from the group consisting of a SAMP6 mouse and a SAMR1 mouse.

10

5

14. The method of claim 11, wherein said determination of apoptotic cells is selected from the group consisting of microscopy of stained cells, TUNEL, Hoescht 33258 dye and video image analysis.

15

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5






FIG. 6



* p<0.05 by ANOVA

FIG. 7



* p<0.05 by ANOVA

FIG. 8





FIG. 9



DBA: dibutyryl cAMP

* p<0.05 by ANOVA

FIG. 10




FIG. 11A




FIG. 11B




FIG. 12




FIG. 13A




FIG. 13B




FIG. 13C

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/23393

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 37/18; A61K 49/00; C12N 5/00, 5/02

US CL : 514/2; 424/9.1; 43/325

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2; 424/9.1; 43/325

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, STN, BIOSIS, CAPLUS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HILL et al. Multiple Extracellular Signals Promote Osteoblast Survival and Apoptosis. Endocrinology. September 1997, Vol.138, No.9, pages 3849-3858.	6,9,10
A	MENG et al. Temporal Expression of the Anabolic Action of PTH in Cancellous Bone of Ovariectomized Rats. Journal of Bone and Mineral Research. 1996, Vol.11, No.4, pages 421-429.	1-5,7,8,11-14
A	CORNISH et al. An in vivo Model for the Rapid Assessment of the Local Effects of Parathyroid Hormone on Bone Histomorphometry. Bone. October, 1995, Vol.17, No.4, supplement, pages 249s-254s.	1-5,7,8,11-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 DECEMBER 1999

Date of mailing of the international search report

11 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KRIS PELHAM MAYO

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/23393

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GUNNESS et al. Anabolic Effect of Parathyroid Hormone on Cancellous and Cortical Bone Histology. Bone. 1993, Vol.14, No.3, pages 277-281.	1-5,7,8,11-14
A	OKAMOTO et al. Femoral Peak Bone Mass and Osteoclast Number in an Animal Model of Age-Related Spontaneous Osteopenia. Anatomical Record. May 1995, Vol.242, No.1, pages 21-28.	1-5,7,8,11-14
A	LECKA-CZERNIK et al. Cellular and Molecular Biomarkers Indicate Precocious in vitro Senescence in Fibroblasts from SAMP6 Mice. Journal of Gerontology. 1997, Vol.52A, No.6, pages B331-B336.	1-5,7,8,11-14
A	JILKA et al. Linkage of Decreased Bone Mass with Impaired Osteoblastogenesis in a Murine Model of Accelerated Senescence. Journal of Clinical Investigation. April 1996, Vol.97, no.7, pages 1732-1740.	1-5,7,8,11-14
A	FUJIBAYASHI et al. Differential Aging Pattern of Cerebral Accumulation of Radiolabeled Glucose and Amino Acid in the Senescence Accelerated Mouse (SAM), a New Model for the Study of Memory Impairment. Biological and Pharmaceutical Bulletin. January 1994, Vol.17, No.1, pages 102-105.	1-5,7,8,11-14